

Final Report for LBA-ECO Team CD-08 Phase I**NCC5-336****Carbon Dynamics in Vegetation and Soils****Susan Trumbore, Jeffrey Q. Chambers****Dept. of Earth System Science, University of California, Irvine, USA****Plinio Camargo, Luiz Martinelli****Lab. Ecologia Isotopica, Centro de Energia Nuclear na Agricultura (CENA),
Piracicaba, Brazil****Joaquim Santos****Instituto de Pesquisa Amazonicas (INPA), Manaus, Brazil****PROJECT SUMMARY**


The overall goals of CD-08 team in Phase I were to quantify the contributions of different components of the carbon cycle to overall ecosystem carbon balance in Amazonian tropical forests and to undertake process studies at a number of sites along the eastern LBA transect to understand how and why these fluxes vary with site, season, and year. We divided this work into a number of specific tasks: (1) determining the average rate (and variability) of tree growth over the past 3 decades; (2) determining age demographics of tree populations, using radiocarbon to determine tree age; (3) assessing the rate of production and decomposition of dead wood debris; (4) determining turnover rates for organic matter in soils and the mean age of C respired from soil using radiocarbon measurements; and (5) comparing our results with models and constructing models to predict the potential of tropical forests to function as sources or sinks of C.

This report summarizes the considerable progress made towards our original goals, which have led to increased understanding of the potential for central Amazon forests to act as sources or sinks of carbon with altered productivity. The overall picture of tropical forest C dynamics emerging from our Phase I studies suggests that the fraction of gross primary production allocated to growth in these forests is only 25-30%, as opposed to the 50% assumed by many ecosystem models. Consequent slow tree growth rates mean greater mean tree age for a given diameter, as reflected in our measurements and models of tree age. Radiocarbon measurements in leaf and root litter suggest that carbon stays in living tree biomass for several years up to a decade before being added to soils, where decomposition is rapid. The time lags predicted from ^{14}C , when coupled with climate variation on similar time scales, can lead to significant interannual variation in net ecosystem C exchange.

B. Background and Objectives

A major goal of the LBA experiment is to determine the status of the Amazon region as a

Figure 7. Carbon cycling in terrestrial ecosystems



source or sink of carbon. Carbon storage capacity in Amazonian Forests, and the capacity for interannual variation in net ecosystem exchange (NEE), is determined by the amount and the residence time of C in ecosystem components. Carbon enters terrestrial ecosystems through a single process, photosynthesis, but can be returned through a variety of processes, collectively referred to as respiration (Figure 1). Net ecosystem exchange can be directly measured by eddy covariance. However, these


measurements do not provide direct information about the processes contributing to the net flux, or why the site may be accumulating or losing carbon (Keller et al., 1996)

The time carbon spends in the ecosystem influences its storage capacity, since it determines how long C stocks can build up before increased decomposition and respiration offset increased productivity. The time elapsed between C uptake by photosynthesis and loss by respiration varies according to how plant C is allocated (to growth or to autotrophic respiration) and the fate of dead plant tissues during decomposition. The residence time of C in tropical forests has been a major unknown limiting our ability to predict how these forests can process and store C. Recently, innovative methods have been developed that use radiocarbon as a direct estimate of the time elapsed since C was removed from the atmosphere by photosynthesis (Trumbore, 2000; Gaudinski et al., 2000; Wang et al., 2000). Our measurements of wood dynamics in permanent plots can be used to estimate the residence time of C in the wood debris pool (Chambers et al., 2001a). The radiocarbon content of leaf and root litter, and soil organic matter may be used to determine the residence time of C in these ecosystem components (Telles, et al., in preparation; Costa et al., in preparation). Finally, the ^{14}C content in CO_2 respired from the soil surface and in incubations may be used to estimate the mean age of C respired from the ecosystem.

Summary of Radiocarbon Tracer

Radiocarbon (^{14}C) is a useful tool for studying the dynamics of C exchange between ecosystems and the atmosphere on several timescales (Trumbore, 2000 and references therein). Radiocarbon is naturally produced by the interaction of high-energy cosmic particles with the upper atmosphere. The ^{14}C formed quickly oxidizes to CO_2 and enters the earth's carbon cycle. The residence time of C in reservoirs that exchange with the atmosphere on century to millennial timescales is determined from the degree to which its ^{14}C has been decreased below atmospheric $^{14}\text{CO}_2$ values by radioactive decay (half-life = 5730 years).

Figure 8. Record of radiocarbon in the atmosphere



Radiocarbon can also be used to estimate C exchange rates on decadal timescales. Atmospheric thermonuclear weapons testing in the 1950's to 1960's roughly doubled the amount of ^{14}C in atmospheric CO_2 in the northern hemisphere prior to the implementation of the Limited Test Ban Treaty in 1963. The rate of incorporation of this 'bomb' ^{14}C provides a measure of the rate of carbon exchange between

atmosphere, ocean, and terrestrial carbon reservoirs on timescales of years to centuries. Models using radiocarbon to determine the turnover time of soil organic matter fractions are summarized in Gaudinski et al (2000) and Trumbore (2000).

Summary of Phase I Activities

To date, we have focused on measurements of C stocks and fluxes at primary forest sites, with emphasis on identifying the major processes controlling C fluxes and dynamics. Measurements include:

- (1) determining the stocks of C (and N) above- and below ground;
- (2) age structure of forest trees and litter components (leaves, branches, roots);
- (3) the influence of climate (seasonal and interannual) and topography on the growth rate of trees;
- (4) the influence of soil texture on storage and dynamics of soil organic matter;
- (5) quantifying fluxes and sources of soil respiration;
- (6) construction of process models to extrapolate in space and time.

Summary of Important Scientific Results

The overall picture of tropical forest C dynamics emerging from our Phase I studies suggests that the fraction of gross primary production allocated to growth in these forests is only 25-30%, as opposed to the 50% assumed by many ecosystem models. Consequent slow tree growth rates mean greater mean tree age for a given diameter, as reflected in our measurements and models of tree age. Radiocarbon measurements in leaf and root litter suggest that carbon stays in living tree biomass for several years up to a decade before being added to soils, where decomposition is rapid. The time lags predicted from ^{14}C , when coupled with climate variation on similar time scales, can lead to significant interannual variation in net ecosystem C exchange.

1. *Predicting the capacity of a central Amazon forest to store carbon. (Chambers et al., 1998; 2000; 2001a,b,c).*




Figure 1. Carbon storage in large wood mass predicted by the stochastic-empirical model developed by Chambers et al (2001a). Changes with simulated ecosystem disturbance (variations about the mean in the first 1000 years), and response to a 25% increase in NPP (ramped up during the period shown by the gray vertical bar) are shown. Rates of C storage are of order 0.5 to 1.0 Mg C ha⁻¹ yr⁻¹ for disturbance.

1.a. Wood dynamics. Chambers et al (1998) constructed a stand dynamics model that integrated 25 years of information on stand dynamics (recruitment, age, and mortality) from permanent forest inventory plots, tree radiocarbon ages, and coarse litter production and decomposition (Chambers et al., 2000). The model simulates the carbon balance of large wood from the dynamics of individual trees. Chambers (2001a) used this model to explore how large wood carbon balance would respond to a 25% increase in net primary production (NPP). Because the model predicts a mean tree age at death of about 175 years, there was a

lag-time of more than a century before large wood standing stock established a new dynamic equilibrium. The predicted annual sequestration rate of about 0.5 Mg C ha⁻¹ yr⁻¹, however, was an order of magnitude less than indicated by a Central Amazon eddy covariance study (Malhi et al. 1998). It is notable that the mean residence time of C in a tree in Chambers' model (~80 years; a 175-year-old-tree is made up of tissues <175 years old) is substantially greater than that used in the CASA model (~40 years). This suggests that (as for soils) lack of inclusion of functional pools with different dynamics oversimplifies the C cycle in non-steady state systems.




Figure 2. Radiocarbon (expressed as the deviation in per mil between the $^{14}\text{C}/^{12}\text{C}$ ratio of the sample and a standard). Positive values indicate the C was fixed from the atmosphere since atomic weapons testing in the 1960s; large negative values indicate that on average C has resided in the soil for centuries (up to -100 per mil) to thousands of years. Except for the Spodosol (an intermittently flooded, sandy, soil sampled in topographic lowland), clay rich Oxisols are contain mostly old carbon at depth.

cycling components reach steady state within about two decades (compared to a century for wood). Comparison of carbon contents of Oxisols sampled near Manaus, Brazil, over the past 20 years shows no measurable change in organic carbon stocks with time, supporting the idea that soils are not currently sequestering a large fraction of ecosystem NPP (Telles et al, in preparation).

2. *Slow growth and great ages for some forest trees (Vieira and Camargo)*

Building on pre-LBA results by Chambers et al. (1998), we have measured radiocarbon age using both direct ^{14}C dating of tree centers and extrapolation from the last 50-years' growth rate. These results (mostly from Manaus and Acre sites) confirm that some small-diameter trees can have ages of several hundred years. Long-term permanent plot studies show some trees with no measurable increase in diameter in 20 years; the ^{14}C method may be the best way of determining growth rates in these trees.




Figure 3. Summary of tree ages estimated from radiocarbon for sites near Manaus (dark circles; Chambers et al, 1998), triangles; new data based on growth rate extrapolations; and Rio Branco (light circles).

1b. Soil C dynamics. Although soil carbon reservoirs are large in deep tropical soils, radiocarbon measurements indicate that the majority of C below ~20 cm depth is refractory (Figure 2, from Telles et al., in preparation). Simple models using radiocarbon and ^{13}C to determine the residence time of C in roots and labile and refractory soil organic C pools indicate that soil C storage rates of 0.5 to .6 Mg C ha⁻¹ yr⁻¹ are possible in the first years following a 25% increase in productivity. Rates of C storage decline rapidly, however, as rapidly

3. Structure and growth rate differences among three primary forests

Biomass. We initiated permanent plot studies at the FLONA km 67 site (in collaboration with CD-10), the Reserva Duke near Manaus (with Higuchi at INPA), and the Catuaba Reserve located near Rio Branco, Acre. Stand dynamics (i.e. recruitment, mortality, damage, and growth) are being monitored annually to determine large-scale carbon balance at these sites. A subset of the trees in each stand are being monitored for monthly growth increment using dendrometer bands; a randomly selected subset of these have been sampled for radiocarbon analyses of decadal growth rates.




Figure 4. Differences in biomass by diameter class.


Estimates of biomass derived using a single allometry equation (Chambers et al., 2000) for the three different sites show dramatic differences between the Central Amazon forest in Manaus, and those in Acre and in the FLONA (Figure 4). Although total biomass is similar, more of the

biomass is in large trees (>60 cm diameter) in the FLONA and Rio Branco sites, while the Manaus site has a greater proportion of its biomass in smaller-diameter (20 cm – 50 cm DBH) trees.

Growth Rates. Preliminary results comparing one year of growth from dendrometer bands in Santarém, Manaus, and Acre permanent plots show much slower growth in trees in Acre. Growth rates do not appear to vary systematically with tree diameter, except for trees <50 cm. We are still analyzing data from Santarém and Acre for seasonal trends.

Our longest-running dendrometer project (Manaus, ZF2), using approximately 300 trees randomly selected from three size classes equally distributed among plateau, slope, and valley forest, was developed as an INPA Forest Science Masters degree project by R. da Silva with Higuchi as advisor and Chambers as co-advisor. Initial results demonstrated significant seasonal variability in diameter increment (Fig. 5), but no variation with

topography (Silva et al. in press). Although there is an obvious relation with season, the relationships between growth rate and



precipitation variables are complex.

Figure 5. Average monthly growth rate and 95% CIs for ~300 trees from DS2. Growth rates increased during the dry to wet season transition months (Sep-Dec), declined throughout the wet season (Jan-May) and were low during the dry season (Jun-August).

4. *A respiration budget for primary tropical forest (Chambers et al., submitted)*

Chambers et al (submitted) extrapolated measurements of soil, leaf, tree bole, and dead tree bole (Chambers et al., 2000;2001a) respiration to estimate the total ecosystem respiration for the ZF2 forest. Soil respiration varied with season and topographic position, with lowest rates in lowland, intermittently inundated soils (*baixios*). Variance in bole respiration was best explained by tree basal diameter and growth rate (from dendrometer measurements). Chambers et al. estimated from their budget that approximately 72% of gross primary production is autotrophically respired. Comparison of “bottom-up” estimates of ecosystem respiration with the tower-based eddy covariance fluxes at night show agreement only on windy nights. While uncertainties in extrapolating leaf-and bole-respiration fluxes to the canopy are large, the study by Chambers et al., (submitted) suggests that respiration and growth largely balance photosynthesis in this central Amazon forest.

5. *There is a significant lag time of several years between photosynthesis and heterotrophic respiration, with much of the lag taking place in living plant tissues (Costa et al., in preparation.)*

The radiocarbon signature of fresh photosynthetic products should equal that of atmospheric CO₂. The ¹⁴C content of atmospheric CO₂ is declining at a rate of 4 to 8 per mil per year, just greater than our measurement accuracy (see section B.1, below). CO₂ measured in soil respiration and evolved during incubation of soils has $\Delta^{14}\text{C}$ values that are 10 to 30 per mil higher than the current years' atmospheric CO₂ (Figure 6), indication that the carbon being respired is on average 2- 4 years old. Radiocarbon measurements of CO₂




Figure 6. $\Delta^{14}\text{C}$ of carbon in fine (<1mm) roots picked from soils in FLONA sites. Although there is large heterogeneity, both live and dead roots have ¹⁴C values consistent with C fixed from the

being respired from soil show large spatial variability, which we hypothesize is linked to leaf litter moisture.

Decomposition in tropical forests occurs rapidly, with leaves decomposing in <1 year and 50% of fine roots in ~1.5 years. The radiocarbon content of root (see below) and leaf litter suggests that the C resides in the living plant tissues for several years (up to decades) before it becomes available for microbial decay (Costa et al., in preparation). Hence a substantial time lag between C uptake by the forest through photosynthesis

and release by decomposition exists in tropical forests that may influence predictions of C storage and release in response to regional variations in climate.

Reasons for the high ^{14}C values of leaves and roots (i.e. true longevity of these components versus the use of reallocated C to produce them) make up part of the PhD thesis of Enir Salazar da Costa and will be emphasized in Phase II measurements.

6. Soil texture plays a major role Carbon cycling in wetter soils differs significantly from upland soils

Clay soils store more carbon than sandy soils along topographic gradients in Santarem and Manaus site (Telles et al., in preparation). However, the extra carbon stored in clay soils is very low in ^{14}C (negative $\Delta^{14}\text{C}$ values; see Figure 2). Highest $\Delta^{14}\text{C}$ values (positive values in Figure 2, meaning the majority of the C was fixed from the atmosphere in the past 50 years) are within 20 cm of the surface in these soils. While ^{14}C values decline rapidly with depth (with greatest decrease in soils with the most clay), the soil with high sand and very low clay content (Manaus Spodosol) maintains high ^{14}C values (the profile stops at the water table), indicating storage of young carbon in this intermittently inundated soil. We intend to increase observations of dissolved organic carbon origins and investigate C cycling in these riparian soils more intensively in Phase II.

Publications during Phase I

- Chambers, J.Q., N. Higuchi, J.P. Schimel, L.V. Ferreira, J. Melack. 2000. Decomposition and carbon cycling of dead trees in tropical evergreen forests of the Central Amazon. *Oecologia* 12:380-388.
- Chambers, J. Q., N. Higuchi, E. S. Tribuzy, and S. E. Trumbore. 2001a. Carbon sink for a century. *Nature* 410: 429.
- Chambers, J. Q., J. P. Schimel, and A. D. Nobre. 2001b. Respiration from coarse wood litter in central Amazon forests. *Biogeochemistry* 52: 115-131.
- Chambers, J.Q., J. Santos, R.J. Ribeiro, and Niro Higuchi. 2001c. Tree damage, allometric relationships, and above-ground net primary production in a tropical forest. *Forest Ecology and Management*. 152:73-84.
- Chambers, J.Q., J. Santos, R.J. Ribeiro, and Niro Higuchi. Submitted. Respiration from a tropical forest ecosystem: An exception to a constant respiration/photosynthesis ratio? *Ecological Applications*.
- Silva, R. P., J. Santos, E. S. Tribuzy, J. Q. Chambers, S. Nakamura, and N. Higuchi. In press. Diameter increment and growth patterns for individual trees in Central Amazon, Brazil. *Forest Ecology and Management*.
- Telles, E.C., S. E Trumbore, P. Camargo, E. Costa, L. A. Martinelli, J. Santos, R. Cosme, Carbon Dynamics and the Capacity for Carbon Storage in Amazonian Forest Soils.

Trumbore, S. Age of soil organic matter and soil respiration: Radiocarbon constraints on belowground C dynamics. *Ecological Applications*, V10(N2):399-411, (2000).

Data Sets Available. Data collected by CD-08 are archived in Brazil and available through the Beija-Flor search engine.

Table 4. Summary of data sets available from CD-08. The final column contains data we need that are being collected by other LBA teams.

MEASUREMENT	LOCATIONS	DATA COLLECTED/PRODUCED	DATA/EQUIPMENT NEEDS
I. Biomass Dynamics			
Permanent plot information	Cautaba Reserve* FLONA* Manaus*	Maps showing locations of plots and individuals; DBH measurements and identification of individuals	
Monthly tree growth rates/wood production	Cautaba Reserve* FLONA* Manaus-Jacaranda*	Monthly diameter increments for trees*, base diameter*, species	precipitation (Higuchi), Monthly PAR (Higuchi), wood density
Annual tree recruitment, growth and mortality; forest structure	Cautaba Reserve* FLONA* Manaus-Ducke	Inventory of base diameters and species*, re-inventory in 2003 and 2006, biomass distribution in base diameter classes, stem density	Botanical Ids (Hopkins at Manaus-Ducke)
Coarse surface litter decomposition*	BDFFP and BIONTE control plots, Manaus	Annual mass loss rate, wood density, tree base diameter, time decomposing	
Large-scale tree mortality/gap formation rates	Manaus-Ducke/EEST	monthly rates from extensive trail network at Ducke, annual rates from remote sensing, minimum size quantifiable from remote sensing and ground truthing	IKONOS and Quickbird (Nelson)
Radiocarbon in cellulose from trees (data product: extrapolated ages)	Catuaba Reserve Manaus* Santarem	Radiocarbon data from cores taken from randomly collected trees in permanent plot sites	Historical climate data for permanent plot sites
II. Heterotrophic Respiration			
Fine surface litter respiration	Jacaranda	Respiratory flux, moisture content, topography, daily changes in moisture (TDR)	Continuous moisture and temperature data in soils
Coarse surface litter respiration	BDFFP and BIONTE control plots*, Jacaranda	Respiratory flux*, moisture content*, wood density*, topography, daily changes in moisture (TDR), TDR/gravimetric calibration	
Bole respiration	Jacaranda plots	Respiratory flux	

Isotopes in soil respired carbon	Paragominas* Santarem* Manaus* Brasilia	Carbon-13 and carbon-14 content of carbon dioxide from purified from soil respiration and soil CO ₂ .	Soil respiration and soil CO ₂ concentration data for these sites (Aduan, Davidson)
III. Soil carbon inventory and isotopes			
Carbon and Nitrogen inventory	Paragominas* Santarem Manaus	Bulk density, carbon, nitrogen content, fine roots picked from soil, soil desription, other soil physical properties	
Isotopes in soil organic matter	Paragominas* Santarem Manaus	Carbon (¹³ C and ¹⁴ C) and nitrogen (¹⁵ N) isotopes in bulk organic matter	
Isotopes in root tissues	Paragomians* Santarem*	Carbon (¹³ C and ¹⁴ C) in live and dead root tissue; root screens to measure isotopes in freshly grown roots	Root inventory (Silver, Nepstad)
Isotopes in leaf and litter tissues	Manaus Santarem Acre	Carbon (¹³ C and ¹⁴ C) compare to leaf specific area and leaf N content	Help with litterfall collection

Notes: * Data already available on LBA-DIS; (name) indicates agreed collaboration for data and or analysis, (name) indicates potential collaboration.

Training and Education. This project has been directly involved in training of 1 US post-doctoral researcher, 3 Brazilian Ph.D. students, 5 MS Brazilian students, 3 undergraduates and one technician.

III. TRAINING AND EDUCATION

People directly involved in research associated with this project are listed in the Table below, with a brief description of their activity. We have had considerable leverage of funds from CNPq and FAPESP – often the role of the NASA funds has been to pay a student until they receive a fellowship.

Name	Level	Funding Source	Description of activity/progress to date
Jeffrey Q. Chambers (USA)	Postdoctoral researcher (UCI/ INPA)	NASA	Co-investigator (promoted to Researcher as of 2/2002) Manaus/INPA measurements, training of students at Manaus
Enir Salazar da Costa (Brazil)	Ph.D student, UC Irvine	NASA	Enir's thesis work includes the C dynamics of leaves and roots in primary tropical forest and the role of these as sources of soil respiration; in addition she is working explore how land management changes cycling of C and N near Santarem
Everaldo de Carvalho Conceição Telles (Brazil)	PhD student, CENA	FAPESP	Finishing PhD in April 2002. Carbon and nitrogen stocks and isotopic signature in soils from Santarem and Manaus with a focus on spatial variability, topography, and C dynamics in soils of different texture.

Simone Aparecida Vieira (Brazil)	PhD student, CENA	FAPESP	Comparison of tree growth rates and forest age structure in Acre, Manaus, and Santarem using permanent plot, dendrometer, and radiocarbon analyses (thesis research in progress)
Roseana Pereira da Silva	MS degree, INPA, 2001 (Chambers, co-advisor)	CNPq	Thesis title: Padrões de crescimento de árvores que ocorrem em diferentes toposseqüências na região de Manaus. Thesis project: Roseana studied monthly changes in tree growth rates using dendrometer bands for trees in three size classes equally distributed among plateau, slope, and valley forests.
Rosana de Miranda Rocha	MS degree, INPA, 2001 (Chambers, co-advisor)	CNPq	Thesis title: Taxas de recrutamento e mortalidade da floresta de terra-firme da bacia do Rio Cuieiras na região de Manaus. Rosana calculated recruitment, growth and mortality rates for two 20 x 2,500 m permanent inventory plots (5 ha each) from 1996 to 2000. She found that the trees were a net carbon sink of about $0.34 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ over the four year period
Bianca Felix	Technician, Manaus	CNPq LBA Bolsa	Chemical analyses of soil and vegetation samples at INPA
Patrícia da Silva Gomes	Undergraduate degree, 2001, INPA (PIBIC)		Densidade de madeira e taxas de crescimento nas parcelas permanentes da reserva florestal ZF-2. Patrícia studied the relationship between wood density and tree growth rates for 60 species on permanent inventory plots at the Biological Dynamics of Forests Fragments Project of the Smithsonian Institution. She found a negative relationship between average wood density and growth rates for these species.
Elder Campos	Undergraduate, Santarem	CNPq LBA Bolsa	Growth band measurements, links to seasonal climate
Kleber Portilho	Undergraduate, Santarem (Chambers, advisor)	CNPq LBA Bolsa	Litter composition and C/N content at km69
Erika Viera de Miranda	MSc student (in progress), INPA (Chambers, co-advisor)	NASA + CNPq	Erika is studying daily changes in stem diameter using high precision linear gauge sensors. She is relating these changes to daily variability in photosynthetically active radiation, volumetric stem water content, and leaf water potential.

Ligia Cristina Toledo	MSc student (in progress), INPA (Chambers, advisor)	NASA + CNPq	Ligia is studying how respiration from coarse and fine litter responds to seasonal changes in precipitation. Her project includes quantifying daily changes in litter moisture content
Liliane Martins Teixiera	MSc student (in progress), INPA (Chambers, advisor)	NASA + CNPq	Liliane is studying how woody tissue respiration has responded to varying logging intensity at the biomass and nutrient experiment (BIONTE) permanent plots. She will also be quantifying how carbon balance differs for low, medium, and high intensity logging treatments.

Other Training and Education activities include participation in short courses (Chambers helped teach the LBA Field course in Manaus, and teaches at INPA; Trumbore and Camargo held a 2-day short course in Isotopes in Ecology course in at the Univ. Federal do Acre, Rio Branco; Trumbore and Pérez taught a similar course at IVIC, Caracas, Venezuela), seminars (by Trumbore at UFAC, Rio Branco; UnB, Brasilia, IVIC (Venezuela); Camargo at INPA, Manaus, UFAC, Rio Branco; UFPa; Santarem) Telles (INPA, Manaus and UFPa, Santarem) and Vieira (UFAC, Rio Branco, INPA, Manaus). Chambers, Telles and Vieira presented posters at the IGBP meeting in Amsterdam in 2001.

Research Significance

This research specifically addresses LBA science questions CD-Q1 (causes on interannual variation in C storage), CD-Q2 (mortality and recruitment, succession), and CD-Q3a (changes in C cycling in pasture/cropland). The information we gathered on processes and timescales of C exchange and storage in Amazon ecosystems is essential for interpreting eddy covariance data, and for extrapolating information from the stand level across broader regions.

REFERENCES CITED

References in bold were published by CD-08 during Phase I.

Camargo, P, S E Trumbore, LA Martinelli, E A Davidson, D C Nepstad, R Victoria, Carbon Dynamics in Regrowing Forest of Eastern Amazonia, *Global Change Biology* V5(N6):693-702 (1999).

Chambers, J. Q., N. Higuchi, and J. P. Schimel. 1998. Ancient trees in Amazonia. *Nature* 391:135-136.

Chambers, J.Q., and S.E. Trumbore. 1999. An age old problem. *Trends in Plant Science* 4: 385-386.

- Chambers, J.Q., N. Higuchi, J.P. Schimel, L.V. Ferreira, J. Melack. 2000. Decomposition and carbon cycling of dead trees in tropical evergreen forests of the Central Amazon. *Oecologia* 12:380-388.
- Chambers, J. Q., N. Higuchi, E. S. Tribuzy, and S. E. Trumbore. 2001a. Carbon sink for a century. *Nature* 410: 429.
- Chambers, J. Q., J. P. Schimel, and A. D. Nobre. 2001b. Respiration from coarse wood litter in central Amazon forests. *Biogeochemistry* 52: 115-131.
- Chambers, J.Q., J. Santos, R.J. Ribeiro, and Niro Higuchi. 2001c. Tree damage, allometric relationships, and above-ground net primary production in a tropical forest. *Forest Ecology and Management*. 152:73-84.
- Chambers, J.Q., J. Santos, R.J. Ribeiro, and Niro Higuchi. Submitted. Respiration from a tropical forest ecosystem: An exception to a constant respiration/photosynthesis ratio? *Ecological Applications*.
- Davidson, E. A. and S. E. Trumbore. Gas diffusivity and production of CO₂ in deep soils of the eastern Amazon, *Tellus*. 47B: 550-565 (1995).
- França, M.B. & Nelson, B.W. (in prep.). Improved estimates of Amazon transition forest biomass. Keller et al., 1996
- Fung, I., C.B. Field, J.A. Berry, M.V. Thompson and J. T. Randerson, C. M. Malmstrom, P. M. Vitousek, G. J. Collatz, P. J. Sellers, D. A. Randall, A. S. Denning, F. Badeck, and J. John. Carbon 13 exchanges between the atmosphere and biosphere. *Global Biogeochemical Cycles* 4:507-533 (1997).
- Gaudinski, J.B., S. E. Trumbore, E. A. Davidson, and Shuhui Zheng, Soil carbon cycling in a temperate forest: radiocarbon-based estimates of residence times, sequestration rates, and partitioning of fluxes, *Biogeochemistry* v. 51:33-69 (2000). Correction/addition: *Biogeochemistry* v. 52:113-114 (2001).
- Malhi, Y., A. D. Nobre, J. Grace, B. Kruijt, M. G. P. Pereira, A. Culf, and S. Scott. 1998. Carbon dioxide transfer over a Central Amazonian rain forest. *Journal of Geophysical Research* :31,593-31,612
- Nepstad, D. C., P. Moutinho, M. B. Dias-Filho, E. Davidson, G. Cardinot, D. Markewitz, R. Figueiredo, N. Vianna, P. Lefebvre, D. Ray, J. Chambers, L. Sternberg, M. Moreira, J. B. Guerrieros, L. Barros, F. Y. Ishida, E. Belk, K. Schwalbe, and I. Tohlver. In press. The effects of partial throughfall exclusion on canopy processes and biogeochemistry of an Amazon forest. *Journal of Geophysical Research*.
- Reich, P. B., M. B. Walters, and D. S. Ellsworth. 1992. Leaf life-span in relation to leaf, plant, and stand characteristics among diverse ecosystems. *Ecological Monographs* 62:365-392.

Schimel, D.S., B.H. Braswell, E.A. Holland, R. McKeown, D.S. Ojima, T.H. Painter, W.J. Parton, and A.R. Townsend. Climatic, edaphic, and biotic controls over storage and turnover of carbon in soils. *Global Biogeochemical Cycles* 8: 279-293 (1994).

Silva, R. P., J. Santos, E. S. Tribuzy, J. Q. Chambers, S. Nakamura, and N. Higuchi. In press. Diameter increment and growth patterns for individual trees in Central Amazon, Brazil. *Forest Ecology and Management*.

Silver, W.L.; Neff, J.; McGroddy, M.; Veldkamp, E.; Keller, M.; Cosme, R. Effects of soil texture on belowground carbon and nutrient storage in a lowland Amazonian forest ecosystem. *Ecosystems*, V3(N2):193-209 (2000).

Telles, E.C., S. E Trumbore, P. Camargo, E. Costa, L. A. Martinelli, J. Santos, R. Cosme, Carbon Dynamics and the Capacity for Carbon Storage in Amazonian Forest Soils.

Trumbore, S. Age of soil organic matter and soil respiration: Radiocarbon constraints on belowground C dynamics. *Ecological Applications*, V10(N2):399-411, (2000).

Vieira, S.; Batista, J.; Selhorst, D.; Brown, F.; Camargo, P.; Martinelli, L.; Trumbore, S. Incertezas na biomassa decorrentes dos inventários - erros de campo (in preparation).

Wang, Y; Amundson, R; Niu, X.F. Seasonal and altitudinal variation in decomposition of soil organic matter inferred from radiocarbon measurements of soil CO₂ flux. *Global Biogeochemical Cycles*, V14(N1):199-211, (2000).

LUIZ ANTONIO MARTINELLI

Birth: 01/23/1959

Associate Professor – University of São Paulo

CENA-USP

Av. Centenário 303,

13416-000, Piracicaba, SP

Brazil

Tel. 55-19-3429.4674

Fax. 55-19-3434-9210

e.mail: zebu@cena.usp.br

Selected Publications (Amazon region only) 1999 - present

Sternberg, L.S.L.; Moreira, M.Z.; Martinelli, L.A.; Victoria, R.L.; Barbosa, E.M.; Bonates, L.C.M.; Nepstad, D.C. The relationship between $^{18}\text{O}/^{16}\text{O}$ and $^{13}\text{C}/^{12}\text{C}$ ratios of ambiente CO_2 in two Amazonian tropical forests. **Tellus** 50: 366-376, 1999.

Medina E, LA Martinelli, E. Barbosa, RL Victoria. Natural abundance of ^{13}C in tropical grasses from INPA herbarium. Revista Brasileira de Botânica 22: 43-51, 1999.

Camargo PB, Trumbore S, Martinelli LA, Davidson EA, Nepstad DC, Victoria RL. Soil carbon dynamics in regrowing forest of eastern Amazonia. **Global Change Biology** 5: 693-702, 1999.

Araújo-Lima CARM, Goulding M, Forsberg BR, Victoria RL, Martinelli LA. The economic value of the Amazonian flooded forest from a fisheries perspective. **Verh. Internat. Verein. Limnol.** 26: 217-2179, 1998

Magnusson WE, Araujo MC, Cintra R, Lima AP, Martinelli LA, Sanaiotii TM, Vasconcelos HL, Victoria RL. Contributions of C-3 and C-4 plants to higher trophic levels in an Amazonian savanna. **Oecologia** 119: 91-96, 1999.

Martinelli LA, Piccolo MC, Townsend AR, Vitousek PM, Cuevas E, McDowell W, Robertson GP, Santos OC, Treseder K. Nitrogen stable isotopic composition of leaves and soil: Tropical versus temperate forests. Biogeochemistry 46: 45-65, 1999.

Martinelli LA, Ballester MV, Krusche A, Victoria RL, Camargo PB, Bernardes M, Ometto JPHB. Land-cover changes and $\delta^{13}\text{C}$ composition of riverine particulate organic matter in the Piracicaba River Basin (southeast region of Brazil). **Limnology and Oceanography** 44: 1826-1833, 1999.

Martinelli LA, Almeida S, Brown IF, Moreira MZ, Victoria RL, Filoso S, Ferreira CAC, and Thomas WW. Variation in Nutrient Distribution and Potential Nutrient losses by Selective Logging in a Humid Tropical Forest of Rondônia, Brazil. **Biotropica** 32: (4) 597-613, 2000

Vitousek PM, Cassman K., Cleveland C, Crews T, Field CB, Grimm NB, Howarth RW, Marino R, Martinelli L, Rastetter EB, Sprent J. Towards an ecological understanding of biological nitrogen fixation. **Biogeochemistry** 48: 1-45, 2000.

Cecilio-Benedito E, Araujo-Lima CARM, Forsberg BR, Bittencourt MM, Martinelli LA. Carbon sources of Amazonian Fisheries. **Fisheries Management and Ecology** 7: 1-10, 2000.

- Magnusson WE, Lima AP, Faria AS, Victoria RL, Martinelli LA. Size and carbon acquisition in lizards from Amazonian savanna: evidence from isotope analysis. **Ecology** 82: 1772-1780, 2001.
- Williams MR, Filoso S, Martinelli LA, Lara LB, Camargo PB. Precipitation and river water chemistry of the Piracicaba River basin, Southeast Brazil. **Journal of Environmental Quality** 30: 967-981, 2001
- Martinelli LA, Camargo PB, Lara LBLS, Victoria RL, Artaxo P. Stable carbon and nitrogen isotopic composition of bulk aerosol particles in a C4 plant landscape of southeast Brazil. **Atmospheric Environment**, in press.
- Sanaïotti T, Martinelli LA, Victoria RL, Trumbore SE, Camargo PB. Past vegetation changes in Amazon savannas by using carbon isotopes of soil organic matter. **Biotropica**, in press.
- Ometto JPHB, Flanagan L, Martinelli LA, Moreira MZ, Higuchi N, Ehleringer J. **Carbon isotope discrimination in forest and pasture ecosystems of the Amazon Basin, Brazil**. *Global Biogeochemical Cycles*, in press.

CURRICULUM VITAE

PLÍNIO BARBOSA DE CAMARGO

Brazilian

18 de julho de 1960

Address: Av. Centenário 303,
13416-000, Piracicaba, SP, Brazil

fone: 55 19 34294788

fax 55 19 34294610

email pcamargo@cena.usp.br

1. Education

Escola Superior de Agricultura "Luiz de Queiroz" - ESALQ/USP-Piracicaba, SP
1981 – 1984, Agonomyst

Centro de Energia Nuclear na Agricultura – CENA/ESALQ/USP-Piracicaba, SP
1985 – 1990, Master, Dissertation: Nitrogen dynamics of the fertilizers uréia (^{15}N) and aquamônia (^{15}N) in the soils with sugar cane.

Centro de Energia Nuclear na Agricultura – CENA/USP-Piracicaba, SP
1992 – 1997, *Doctor, Thesis: Soil carbon organic matter dynamics in the soils of Amazonia because land use change.*

2. Publications

- CAMARGO P.B.de, SALOMÃO R. de P., TRUMBORE S., MARTINELLI L.A.. How old are large Brazil-nut trees (*Bertholletia excelsa*) in the Amazon? *Scientia Agricola*, Piracicaba, 51(2):389-391, maio/ago.1994.

- VICTORIA R.L., FERNANDES F., MARTINELLI L.A., PICCOLO M. de C., CAMARGO P.B.de, TRUMBORE S.. Past vegetation changes in the Brazilian Pantanal arboreal-grassy savanna ecotone by using carbon isotopes in the soil organic matter. *Global Change Biology* 1(3):165-171, 1995.

- TRUMBORE S.E., DAVIDSON E.A., CAMARGO P.B.de, NEPSTAD D.C., MARTINELLI L.A.. Belowground cycling of carbon in forests and pastures of Eastern Amazonia. *Global Biogeochemical Cycles*, 9(4):515-528, december, 1995.

- MARTINELLI L.A., PESSENDA L.C.R., ESPINOZA E.P., CAMARGO P.B.de, TELLES E.C., CERRI C.C., VICTORIA R.L., ARAVENA R., RICHEY J., TRUMBORE S.. Carbon-13 depth variation in soils of Brazil and relations with climate changes during the Quaternary. *Oecologia*, 106:376-381, 1996.

- PESSENDA L.C.R., VALENCIA E.P.E., CAMARGO P.B.de, TELLES E.C.C.; MARTINELLI L.A., CERRI C.C., ARAVENA R., ROZANSKI K.. Natural radiocarbon measurements in Brazilian soils developed on basic rocks. *Radiocarbon*, Tucson, 38(2):203-208, 1996.

- MARTINELLI L.A. & CAMARGO P.B.de. O monitoramento do carbono em áreas florestadas. *Série Técnica IPEF*, Piracicaba, v.10, n.29, p.6-10, 1996.

- KRUSCHE A.V., CARVALHO F.P. de, MORAES J.M. de, CAMARGO P.B. de, BALLESTER M.V.R., HORNINK S., MARTINELLI L.A., VICTORIA R.L.. Spatial and temporal water quality variability in the Piracicaba river basin, Brazil.

Journal of the American Water Resources Association, 33(5), pg.1117-1123, october 1997.

- **MARTINELLI L.A., KRUSCHE A.V., VICTORIA R.L., CAMARGO P.B.de, BERNARDES M.C., FERRAZ E.S., MORAES J.M., BALLESTER M.V.R., Effects of sewage on the chemical composition of Piracicaba river, Brazil. *Water, Air and Soil Pollution*, 110:67-79, 1999**

- CAMARGO P.B.de, TRUMBORE S.E., MARTINELLI L.A., DAVIDSON E.A., NEPSTAD D.C., VICTORIA R.L.. Soil carbon dynamics in regrowing forest of eastern Amazonia. *Global Change Biology*, 5(6). Pg. 693-702, 1999.

- BALLESTER M.V.R., MARTINELLI L.A., KRUSCHE A.V., VICTORIA R.L., BERNARDES M.C., CAMARGO P.B.de. Effects of increasing organic matter loading on the dissolved O₂, free dissolved CO₂ and respiration rates in the Piracicaba river basin, southeast Brazil. *Water Resources*, 33(9), pg.2119-2129, 1999.

- PISSINATO L., MARTINELLI L.A., VICTORIA R.L., CAMARGO P.B.de. Using stable carbon isotopic analyses to access the botanical origin of ethanol in Brazilian brandies. *Food Research International* 32: 665-668, 1999.

- MARTINELLI L.A., BALLESTER M.V., KRUSCHE A., VICTORIA R.L., CAMARGO P.B.de, BERNARDES M., OMETTO J.P.H.B.. Land-cover changes and $\delta^{13}\text{C}$ composition of riverine particulate organic matter in the Piracicaba River Basin (southeast region of Brazil). *Limnology and Oceanography* 44: 1826-1833, 1999.

- PEREZ T., TRUMBORE S.E., TYLER S.C., DAVIDSON E.D, KELLER M., CAMARGO P.B.de. Isotopic variability of N₂O emissions from tropical forest soils. *Global Biogeochemical Cycles*, 14(2), pg.525-535, 2000.

- SILVEIRA A.M., VICTÓRIA R.L., BALLESTER M.V., CAMARGO P.B.de, MARTINELLI L.A., PICCOLO M.C.Simulação dos efeitos das mudanças do uso da terra na dinâmica de carbono no solo na bacia do rio Piracicaba. *Pesquisa Agropecuária Brasileira*, 35(2), pg.389-399, 2000.

- WILLIAMS M.R., FILOSO S., MARTINELLI L.A., LARA L.B., CAMARGO P.B. Precipitation and river water chemistry of the Piracicaba River basin, Southeast Brazil. *Journal of Environmental Quality* 30: 967-981, 2001

- LARA L.B.S.L., ARTAXO P., MARTINELLI L.A., VICTORIA R.L., CAMARGO P.B. DE, KRUSCHE A., AYERS G.P., FERRAZ E.S.B., BALLESTER M.V.. Chemical composition of rainwater and anthropogenic influences in the Piracicaba River basin, southeast Brazil. *Atmospheric Environment* 35: 4937-4945, 2001.

- DANIEL M.H.B., MONTEBELLO A.A., BERNARDES M.C., OMETTO J.P.H.B., CAMARGO P.B., KRUSCHE A.V., BALLESTER M.V., VICTORIA R.L., MARTINELLI L.A.. Effects of urban sewage on dissolved oxygen dissolved inorganic carbon and organic carbon, and electrical conductivity of small streams along a gradient of urbanization in the Piracicaba River basin. *Water, Air, and Soil Pollution*, in press.

- MARTINELLI L.A., CAMARGO P.B., LARA L.B.L.S., VICTÓRIA R.L., ARTAXO P.. Stable carbon and nitrogen isotopic composition of bulk aerosol particles in a C4 plant landscape of southeast Brazil. *Atmospheric Environment*, in press.

- SANAIOTTI T., MARTINELLI L.A., VICTORIA R.L., TRUMBORE S.E., CAMARGO P.B.. Past vegetation changes in Amazon savannas by using carbon isotopes of soil organic matter. *Biotropica*, in press.

3. Congress and Workshops

- CAMARGO P.B.de, TRUMBORE S.E., DAVIDSON E.D., NEPSTAD D.C., MARTINELLI L.A.. $\delta^{14}\text{C}$, $\delta^{13}\text{C}$ and C Inventory Differences Between Forest and Pasture in Soils of Eastern Amazonia. Supplement to Eos, Transactions, American Geophysical Union. 1993 Fall Meeting. Vol.74, N°43, outubro de 1993.

- CAMARGO P.B.de, MARTINELLI L.A., PESSENDA L.C.R., TRUMBORE S.E., NEPSTAD D.C., DAVIDSON E.D.. Evaluation of C and N inventories, $\delta^{13}\text{C}$ e $\Delta^{14}\text{C}$ in deep soils from mature, regrowing forest (capoeira) and pasture of Eastern Amazonia. Abstracts of: Eighth International Conference on Geochronology Cosmochronology and Isotope Geology. Berkeley, CA, USA, pg 46, 5 a 11 de junho de 1994.

- CAMARGO P.B.de, MARTINELLI L.A., TRUMBORE S.E., NEPSTAD D.C.. Soil carbon cycling in deep soil from forests and pastures of Eastern Amazonia. In Abstracts of: XII International Symposium on Environmental Biogeochemistry. Biosphere and Atmospheric Changes. Rio de Janeiro, RJ, Brazil, pg 61, 3 a 8 de setembro de 1995.

- CAMARGO P.B.de, TRUMBORE S.E., VIEIRA S.A., CHAMBERS J.A., HIGUCHI N., MARTINELLI L.A., SOUTON J.. Radial growth rates of trees with different wood densities in eastern Amazonia determined from radiocarbon American Geophysical Union 81(48), pg. F185, 2000.

- TELLES E.C., CAMARGO P.B.de, TRUMBORE S.E., MARTINELLI L.A., VICTORIA R.L., MOREIRA M.Z., SANTOS J.. Spatial variation and correlation of ^{13}C , ^{15}N , C and N content in Oxisols and Ultisols in primary forest of Amazonia. American Geophysical Union, 81(48), pg.F186, 2000.

- TRUMBORE S.E., CAMARGO P.B.de, NEPSTAD D.C., MARKEWITZ D., DAVIDSON E., FIGUEIREDO R., ISHIDA F.Y.. Age of fine roots in terra firme forests of eastern Amazonia and the influence of roots on the ^{14}C of soil respiration. American Geophysical Union, 81(48), pg.F234, 2000.

- TELLES E.C., CAMARGO P.B.de, TRUMBORE S.T., MARTINELLI L.A., VICTORIA R.L., MOREIRA M.Z., SANTOS J.. Prediction of delta ^{13}C for yellow latosols of amazon under primary forest. First LBA Scientific Conference, pg 121, 2000.

- VIEIRA S.A., CAMARGO P.B.de, TRUMBORE S.E., BROWN I.F., MENDOZA E.R.H., SELHORST D., CHAMBERS J.C., SANTOS J.. Mudanças globais e taxa de crescimento das espécies arbóreas da Amazonia. First LBA Scientific Conference, pg. 128, 2000

- PÉREZ T, TRUMBORE S.E., TYLER S.C., DAVIDSON E.A., KELLER M., CAMARGO P.B.de.. Isotopic variability of N_2O emissions from tropical forest: monitoring soil texture impact and spacial variability. First LBA Scientific Conference, 76, 2000.

Dr. Plínio Barbosa de Camargo
Piracicaba, March 03, 2002